

Neutron Scattering and d -Wave Symmetry in the Cuprates

In this comment we demonstrate that the general conjectures of Fong *et al* on their $\text{YBa}_2\text{Cu}_3\text{O}_7$ neutron experiments are realized in concrete calculations of the neutron scattering structure factor $S(\mathbf{q}, \omega)$ in this material. We make predictions for similar experiments in $\text{La}_2\text{Sr}_{0.15}\text{Cu}_{1.85}\text{O}_4$, which yield a different q -structure due to the different fermiology. Our calculations address both the frequency and wave-vector dependence of the data and (1) include realistic Fermi surface effects (2) are compatible with the normal state data and (3) demonstrate the consistency of a d -wave order parameter, in $\text{YBa}_2\text{Cu}_3\text{O}_7$ although they appear to be also compatible with an alternate symmetry specific to the orthorhombic bi-layer system.

We have shown [2] that the normal state neutron data in both $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$ and $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ can be understood as a competition between fermiology and antiferromagnetic exchange effects $J(\mathbf{q})$. An infinite Coulomb U expansion leads to an RPA-like theory which yields an $S(\mathbf{q}, \omega)$ with wave-vector, $\mathbf{q} = (q_x, q_y)$ [as well as q_z in the bi-layer system] and ω dependences observed experimentally. Similarly, a four peak signature of the d -wave state at very low ω was presented, although experiments on both cuprates have failed to confirm this latter d -wave behavior.

Using our theory we plot the normal and superconducting data for $\text{YBa}_2\text{Cu}_3\text{O}_7$ in Figure 1a as a function of $\mathbf{q} = (q_x, q_y)$ at frequencies $\omega = 41\text{meV}$ slightly larger than twice the d -wave gap frequency $2\Delta_o$. The results are in reasonable agreement with Ref 1, although the calculated width is somewhat wider than observed. The dependence on $\mathbf{q} = (q_x, q_y)$ is similar for both $q_z = 0$ and $q_z = \pi$, but the amplitude is only significant for the latter wave vector [3]. This follows in large part because the two d -wave gaps in each of the bilayer sub-bands are not appreciably different [4] on a scale of 41 meV. The mechanism for the formation of the strong peak below T_c is, as conjectured in Ref 1, a consequence of pair creation with one electron in the + and the other electron in the - lobe of the order parameter, whose wave vector separation is near $(q_x, q_y) = (\pi, \pi)$. This enhances the weight around $(q_x, q_y) = (\pi, \pi)$ in the Lindhard function. The further inclusion of an antiferromagnetic exchange $J(\mathbf{q})$ creates a strong peak at $(q_x, q_y) = (\pi, \pi)$. Comparable calculations are presented in Figure 1b for $\text{La}_2\text{Sr}_{0.15}\text{Cu}_{1.85}\text{O}_4$, where the Fermi surface nesting should lead to incommensurate d -wave peaks. Here the characteristic frequency is taken to be 17 meV slightly larger than 16 meV for the estimated $2\Delta_o$. While experiments have been limited to lower ω , future data should focus on this regime.

We end by noting that for the bi-layer system, the presence of orthorhombicity can create a state [4] with a +s gap on one band and a -s gap on the other, each

of which gap is oriented along a different in-plane crystal axis. This mimics a d -wave state and should also be compatible with the neutron data of Ref. 1. A similar state, however, has no analogue in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$.

This work was supported by NSF-DMR-91-20000 and NSF-DMR-94-16926.


Yuyao Zha^a and K. Levin^b

^aDepartment of Physics, University of Illinois, 1110 W. Green Street, Urbana, IL 61801

^bThe James Franck Institute, The University of Chicago 5640 South Ellis Avenue, Chicago, IL 60637

FIG. 1. Calculated neutron scattering form factor $\text{Im}\chi$ as a function of (q_x, q_y) according to the model described in Ref [2], for (a) $\text{YBa}_2\text{Cu}_3\text{O}_7$ by assuming the maximum of the superconducting gap $\Delta_o = 20\text{meV}$, $J_0/J_c = 0.7$; and (b) $\text{La}_2\text{Sr}_{0.15}\text{Cu}_{1.85}\text{O}_4$ for $\Delta_o = 8\text{meV}$, $J_0/J_c = 0.6$.

-
- [1] Hung Fai Fong *et al*, preprint.
 - [2] Y. Zha *et al*, Phys. Rev. B 47, 9124(1993); Q. Si *et al*, Phys. Rev. B 47, 9055(1993).
 - [3] Y. Zha *et al*, Physica C 212, 413-418(1993).
 - [4] D. Z. Liu, K. Levin and J. Maly, preprint.



$La_{1.85}Sr_{0.15}CuO_4$

(b)

$\omega=17meV$

